POST-TEST

Year in Review: Clinical Investigator Perspectives on the Most Relevant New Datasets and Advances in Colorectal Cancer

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. The Phase III AZUR-2 trial is investigating the efficacy and safety of which of the following agents in the perioperative setting for patients with Stage III microsatellite instability-high/mismatch repairdeficient (dMMR) resectable colorectal cancer (CRC)?
 - a. Nivolumab
 - b. Pembrolizumab
 - c. Dostarlimab
- 2. Which of the following statements best represents outcomes of the Phase III CALGB/SWOG-80702 trial evaluating the effect of celecoxib versus placebo added to standard adjuvant therapy for patients with Stage III CRC?
 - No significant improvement in disease-free survival (DFS) for patients with or without detectable circulating tumor DNA (ctDNA)
 - b. Significant improvement in DFS for patients with detectable ctDNA
 - c. Significant improvement in DFS for patients with or without detectable ctDNA

3. Which of the following statements best characterizes a clinical research objective of the ongoing Phase II/III NRG-GI008 trial?

- a. To compare DFS rates in the cohort of patients without detectable ctDNA after receiving first-line FOLFOX versus FOLFOX/bevacizumab
- b. To compare DFS rates in the cohort of patients with detectable ctDNA after receiving first-line FOLFOX versus FOLFOX/bevacizumab
- c. To compare DFS rates in the cohort of patients without detectable ctDNA after disease resection and receiving immediate versus delayed chemotherapy

- 4. What was the approximate clinical complete response rate in a trial evaluating neoadjuvant dostarlimab for Stage II/III rectal cancer?
 - a. 0%
 - b. 25%
 - c. 50%
 - d. 75%
 - e. 100%
- 5. The rate of pathologic response observed in the NICHE-2 trial of neoadjuvant nivolumab with ipilimumab for locally advanced dMMR colon cancer is best characterized by which of the following statements?
 - a. No pathologic responses were observed in patients
 - b. A pathologic response was observed in a low proportion of patients
 - c. A pathologic response was observed in a high proportion of patients