



POST-SABCS Issue 2, 2013

Confirmatory Overall Survival Analysis of CLEOPATRA

For more visit ResearchToPractice.com/5MJCSABCS2013

Research
To Practice®

CME INFORMATION

OVERVIEW OF ACTIVITY

The annual San Antonio Breast Cancer Symposium (SABCS) is unmatched in its significance with regard to the advancement of breast cancer treatment. This unique conference provides many clinicians and scientists from around the world a forum for discussing critical issues in the prevention and management of breast cancer. Therefore, SABCS is targeted by many members of the clinical research community as the optimal time to unveil new clinical data. This creates an environment in which yearly published results from a plethora of ongoing clinical trials lead to the emergence of new therapeutic agents and changes in the indications for existing treatments across all breast cancer subtypes.

Although SABCS currently offers online access to the vast majority of the poster and plenary presentations from the annual meeting, the absence of expert assessment concerning practical applications may yield confusion among community oncologists who are challenged almost daily to appropriately apply a multitude of scientific findings across diverse tumors. Thus, identification of those data sets with immediate or impending relevance to the delivery of quality breast cancer care, in conjunction with professional commentary to address resultant practice ambiguity, may prove vastly beneficial to those physicians who are unable to make the annual pilgrimage to San Antonio. To bridge the gap between research and patient care, this CME activity will deliver a serial review of the most important emerging data sets from the latest SABCS, including expert perspectives on how these new evidence-based concepts can and should be incorporated into on- and off-protocol patient care. This activity will assist medical oncologists, radiation oncologists, breast/general surgeons, nurses and other healthcare professionals in the formulation of optimal clinical management strategies and the timely application of new research findings to best-practice patient care.

LEARNING OBJECTIVES

- Appraise recent clinical research findings from the second interim survival analysis of the CLEOPATRA study and the subset analysis of patients based on age, and apply this information to the treatment of patients with metastatic HER2-positive breast cancer.
- Recall the benefits and risks of combining HER2-targeted antibodies with chemotherapeutic agents for the treatment of HER2-positive advanced breast cancer.
- Understand the association between PI3 kinase mutational status and prognosis in patients with HER2-positive metastatic breast cancer.
- Evaluate the efficacy and safety of adding eribulin mesylate to trastuzumab for patients with HER2-positive advanced breast cancer.
- Compare the toxicity profile of T-DM1 across multiple studies in metastatic HER2-positive breast cancer, and consider this information in the selection of optimal HER2-targeted later-line therapy.

ACCREDITATION STATEMENT

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

CREDIT DESIGNATION STATEMENT

Research To Practice designates this enduring material for a maximum of 1.75 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

HOW TO USE THIS CME ACTIVITY

This CME activity contains slides and edited commentary. To receive credit, the participant should review the slide presentations, read the commentary, complete the Post-test with a score of 75% or better and fill out the Educational Assessment and Credit Form located at ResearchToPractice.com/5MJCSABCS2013/2/CME.

CONTENT VALIDATION AND DISCLOSURES

Research To Practice (RTP) is committed to providing its participants with high-quality, unbiased and state-of-the-art education. We assess potential conflicts of interest with faculty, planners and managers of CME activities. Real or apparent conflicts of interest are identified and resolved through a conflict of interest resolution process. In addition, all activity content is reviewed by both a member of the RTP scientific staff and an external, independent physician reviewer for fair balance, scientific objectivity of studies referenced and patient care recommendations.

FACULTY — The following faculty (and their spouses/partners) reported real or apparent conflicts of interest, which have been resolved through a conflict of interest resolution process:

José Baselga, MD, PhD
Physician-in-Chief
Memorial Sloan-Kettering Cancer Center
New York, New York

Consulting Agreements: Genentech BioOncology, Novartis Pharmaceuticals Corporation.

Kimberly L Blackwell, MD
Professor of Medicine
Director, Breast Cancer Program
Duke Cancer Institute
Durham, North Carolina

Advisory Committee: Novartis Pharmaceuticals Corporation;
Consulting Agreements: Novartis Pharmaceuticals Corporation, Sandoz; Contracted Research: Celgene Corporation, Genentech BioOncology, Roche Laboratories Inc; Speakers Bureau: Genomic Health Inc.

Lisa A Carey, MD
Richardson and Marilyn Jacobs Preyer Distinguished Professor for Breast Cancer Research
Chief, Division of Hematology and Oncology
Physician-in-Chief
North Carolina Cancer Hospital
Associate Director for Clinical Research
Lineberger Comprehensive Cancer Center
Chapel Hill, North Carolina

Advisory Committee, Consulting Agreements and Speakers Bureau: Amgen Inc, Bristol-Myers Squibb Company, Genentech BioOncology, Novartis Pharmaceuticals Corporation, Pfizer Inc, Sanofi; Research Support: Genentech BioOncology, GlaxoSmithKline, Sanofi.

Edith A Perez, MD
Deputy Director at Large, Mayo Clinic Cancer Center
Group Vice Chair, Alliance of Clinical Trials in Oncology
Serene M and Frances C Durling Professor of Medicine
Mayo Clinic
Jacksonville, Florida

Contracted Research: Genentech BioOncology, Genomic Health Inc, GlaxoSmithKline.

EDITOR — Dr Love is president and CEO of Research To Practice, which receives funds in the form of educational grants to develop CME activities from the following commercial interests: AbbVie Inc, Algeta US, Allos Therapeutics, Amgen Inc, ArQule Inc, Astellas, Aveo Pharmaceuticals, Bayer HealthCare Pharmaceuticals, Biodesix Inc, Biogen Idec, Boehringer Ingelheim Pharmaceuticals Inc, Bristol-Myers Squibb Company, Celgene Corporation, Daiichi Sankyo Inc, Dendreon Corporation, Eisai Inc, EMD Serono Inc, Foundation Medicine Inc, Genentech BioOncology, Genomic Health Inc, Gilead Sciences Inc, Incyte Corporation, Lilly USA LLC, Medivation Inc, Merck, Millennium: The Takeda Oncology Company, Mundipharma International Limited, Novartis Pharmaceuticals Corporation, Onyx Pharmaceuticals Inc, Prometheus Laboratories Inc, Regeneron Pharmaceuticals, Sanofi, Seattle Genetics, Spectrum Pharmaceuticals Inc and Teva Oncology.

RESEARCH TO PRACTICE STAFF AND EXTERNAL REVIEWERS — The scientific staff and reviewers for Research To Practice have no real or apparent conflicts of interest to disclose.

This educational activity contains discussion of published and/or investigational uses of agents that are not indicated by the Food and Drug Administration. Research To Practice does not recommend the use of any agent outside of the labeled indications. Please refer to the official prescribing information for each product for discussion of approved indications, contraindications and warnings. The opinions expressed are those of the presenters and are not to be construed as those of the publisher or grantors.

This activity is supported by educational grants from Eisai Inc, Genentech BioOncology and Genomic Health Inc.

Hardware/Software Requirements:

A high-speed Internet connection

A monitor set to 1280 x 1024 pixels or more

Internet Explorer 7 or later, Firefox 3.0 or later, Chrome, Safari 3.0 or later

Adobe Flash Player 10.2 plug-in or later

Adobe Acrobat Reader

(Optional) Sound card and speakers for audio


Last review date: March 2013

Expiration date: March 2014

The new world of HER2-positive breast cancer

To go directly to slides and commentary for this issue, [click here](#).

Last week I met with the new Physician-in-Chief at Memorial Sloan-Kettering Cancer Center, Dr José Baselga, and after talking a bit about his vision for the future of that preeminent institution we focused on a corner of oncology he has influenced mightily throughout his career, breast cancer research. Not surprisingly, we spent much of our time reviewing anti-HER2 treatment — which has witnessed the FDA approval of 2 new agents in the past 9 months. Dr Baselga got things started by commenting on the Phase III trial he chaired, CLEOPATRA, which clearly demonstrated a substantial boost in efficacy when the HER2 dimerization inhibitor pertuzumab (P) was added to docetaxel and trastuzumab (T) as first-line therapy for metastatic HER2-positive disease. In the trial progression-free survival (PFS) increased from 12.4 to 18.5 months with a similar safety profile, and although the magnitude of this landmark finding surprised many observers, Dr Baselga stated that he fully expected the results based on the substantial antitumor activity seen when P was added to T in patients with disease progression on T in a prior Phase II trial.



José Baselga, MD, PhD
RTP Studios (3-9-2013)

We then chatted about the antibody-drug conjugate trastuzumab emtansine (T-DM1) and the EMILIA trial that exploded onto the scene during the ASCO 2012 plenary session, revealing T-DM1's clear-cut superiority in both efficacy (PFS and overall survival) and tolerability over an established and frequently used regimen (capecitabine/lapatinib) among patients who had previously been treated with T + a taxane. As the agent was just approved 2 weeks before the interview, our conversation took on a different tone, as for the first time I was able to ask an investigator the practical (rather than hypothetical) question of current sequencing of therapy for metastatic HER2-positive disease. Dr Baselga, in commenting on this complex issue that has likely been discussed at every tumor board on the planet, slowly removed his eyeglasses, carefully put them on the desk, thought for a moment and then voiced his perspective, which is similar to

those I have heard recently from Dr Eric Winer and others: "Some people are so excited about T-DM1 that they want to use it first line, but I think this is a time for intellectual calm. Right now, trastuzumab, pertuzumab and a taxane is our standard first-line treatment, with T-DM1 as second line."

For the record, he and his Memorial colleagues usually choose paclitaxel as a partner for T + P, partially based on the reassuring Phase II data the group reported at San Antonio with this regimen. As I have been known to do, I tried to push Dr Baselga a bit regarding his strong feeling not to use T-DM1 first line and asked him how he would approach an 85-year-old patient with ER-negative, HER2-positive metastatic breast cancer for whom traditional chemotherapy might be out of place. He, however, stuck to his guns, commenting that a short taxane course (with T + P) in many fit, older patients is a well-tolerated life investment that results in a median progression-free interval of 18 months.

Whatever the algorithm is for now, it may very well change in a year or so when the MARIANNE study reports. This crucial Phase III first-line trial compares T + a taxane to T-DM1 alone or with P. Dr Baselga very clearly stated his opposition to the nonprotocol use of T-DM1 combined with P until more trial data become available, and other investigators, including Dr Winer, have done the same.

Of course, many other complex questions remain about the treatment of metastatic HER2-positive breast cancer, and below we review some of the more interesting efforts unveiled in San Antonio that attempt to provide needed answers:

1. More from the CLEOPATRA trial: Overall survival benefit; biomarker analysis; effects in older patients

With 154 deaths in the control group and 113 in the T + P + docetaxel arm, [the study](#) has now allowed crossover to P. In terms of biomarkers, according to Dr Baselga, who presented [these data](#) in San Antonio, perhaps the key factor moving forward will be the identification of PI3-kinase mutations in approximately 25% of HER2-positive tumors and the potential use of PI3-kinase alpha inhibitors, which are currently being evaluated. Finally, although only 126 patients in [CLEOPATRA](#) were older than age 65, the benefit they derived from treatment was similar to what was seen with younger patients.

2. Choice of chemotherapy to combine with T + P

Referred to earlier, a [San Antonio report](#) from Memorial demonstrated a 76% 6-month PFS rate in 33 evaluable patients receiving T + P + weekly paclitaxel. No unexpected toxicities were encountered, and this work provides additional strength to the conclusion everyone, including the NCCN, had already reached, namely that paclitaxel is a reasonable agent to combine with T and P. To obtain more real-world perspectives on this issue, an international single-arm study ([PERUSE](#)) is now evaluating T + P with 3 different taxanes (paclitaxel, *nab* paclitaxel and docetaxel).

3. Pooled safety analysis of single-agent T-DM1

These data from 882 patients on 6 clinical trials (including EMILIA) revealed few clinically apparent toxicities but did document transient laboratory abnormalities, such as thrombocytopenia and abnormal liver function tests, in a quarter or more of patients. Overall, treatment discontinuation due to toxicity was observed in only 55 patients (6.2%).

4. Eribulin combined with T

Indefinite anti-HER2 treatment is now a standard part of care for patients with HER2-positive metastatic disease, and as new chemotherapy agents are developed, studies are needed to document whether these are safe and efficacious partners for T. **This report** of 40 patients demonstrated what most observers expected — efficacy similar to other chemotherapy/T combinations (55% CR + PR) and acceptable tolerability comparable to what has been reported with eribulin alone.

For the next issue of this series we review the many San Antonio papers on genomic markers, including yet another analysis with the 21-gene recurrence score in tumor specimens from a prior randomized adjuvant trial, in this case NSABP-B-28, which evaluated the addition of paclitaxel to AC.

Neil Love, MD

Research To Practice

Miami, Florida

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Research To Practice designates this enduring material for a maximum of 1.75 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

This activity is supported by educational grants from Eisai Inc, Genentech BioOncology and Genomic Health Inc.

Confirmatory Overall Survival Analysis of CLEOPATRA

Presentation discussed in this issue

Swain SM et al. **Confirmatory overall survival (OS) analysis of CLEOPATRA: A randomized, double-blind, placebo-controlled Phase III study with pertuzumab (P), trastuzumab (T), and docetaxel (D) in patients (pts) with HER2-positive first-line (1L) metastatic breast cancer (MBC).** San Antonio Breast Cancer Symposium 2012; **Abstract P5-18-26.**

Slides from a presentation at SABCS 2012 and transcribed comments from a recent interview with Edith A Perez, MD (1/17/13)


Confirmatory Overall Survival (OS) Analysis of CLEOPATRA: A Randomized, Double-Blind, Placebo-Controlled Phase III Study with Pertuzumab, Trastuzumab, and Docetaxel in Patients (pts) with HER2-Positive First-Line (1L) Metastatic Breast Cancer (MBC)

Swain SM et al.

Proc SABCS 2012; Abstract P5-18-26.

Research
To Practice®

CLEOPATRA Study Design



Primary endpoint: Independently assessed progression-free survival (PFS)

Baselga J et al. *N Engl J Med* 2012;366(2):109-19.

Research
To Practice®

CLEOPATRA: Primary Efficacy Analysis


Independently assessed	Ptz + T + D (n = 402)	Pla + T + D (n = 406)	Hazard ratio	p-value
Median PFS	18.5 mo	12.4 mo	0.62	<0.001
Deaths in interim overall survival analysis, n (%)*	69 (17.2%)	96 (23.6%)	0.64	0.005

* Not significant because analysis did not meet O'Brien-Fleming stopping boundary; a strong trend was evident toward overall survival benefit with pertuzumab

Baselga J et al. *N Engl J Med* 2012;366(2):109-19.

Research
To Practice®

CLEOPATRA: Overall Survival (Confirmatory Analysis)




- Crossed O'Brien-Fleming stopping boundary and was therefore deemed statistically significant.
- Analysis was performed after 267 deaths and 69% of the prespecified total number of events for the final analysis had occurred.
- Median follow-up in both arms = 30 months

With permission from Swain SM et al. *Proc SABCS 2012*; Abstract P5-18-26.

Research
To Practice®


CLEOPATRA: Overall Survival in Predefined Subgroups



With permission from Swain SM et al. *Proc SABCS 2012*; Abstract P5-18-26.

Research
To Practice®

CLEOPATRA: Follow-Up Analysis — Updated Investigator-Assessed PFS



- At the time of the data cutoff, 296 (72.9%) patients on the placebo arm and 257 (63.9%) on the pertuzumab arm had experienced a PFS event. These results were exploratory only.
- This updated analysis of investigator-assessed PFS was consistent with the results from the primary PFS analyses.

With permission from Swain SM et al. *Proc SABCS 2012*;Abstract P5-18-26.

Research
To Practice®

CLEOPATRA: Select All-Grade Adverse Events (AEs)

AEs with $\geq 25\%$ incidence or $\geq 5\%$ difference between arms	Ptz + T + D (n = 408)	Pla + T + D (n = 396)
Diarrhea	68.1%	48.2%
Rash	36.5%	24.0%
Mucosal inflammation	27.5%	19.9%
Peripheral edema	24.8%	30.8%
Pruritus	16.7%	10.1%
Constipation	15.4%	25.5%
Febrile neutropenia	13.7%	7.6%
Dry skin	10.8%	5.8%

Ptz + T + D did not increase the incidence of cardiac adverse events compared to Pla + T + D.

Swain SM et al. *Proc SABCS 2012*;Abstract P5-18-26.

Research
To Practice®

CLEOPATRA: Grade ≥ 3 Adverse Events

Grade ≥ 3 adverse events (incidence $\geq 5\%$)	Ptz + T + D (n = 408)	Pla + T + D (n = 396)
Neutropenia	49.0%	46.0%
Febrile neutropenia	13.7%	7.6%
Leukopenia	12.3%	14.9%
Diarrhea	9.1%	5.1%

Mucosal inflammation (Grade ≥ 3) was reported in 1% of patients in the placebo arm and 1.5% of patients in the pertuzumab arm.

Swain SM et al. *Proc SABCS 2012*;Abstract P5-18-26.

Research
To Practice®

Author Conclusions

- At the second interim analysis, CLEOPATRA demonstrated a statistically significant and clinically meaningful improvement in OS with the addition of pertuzumab to trastuzumab/docetaxel as first-line therapy for HER2-positive mBC.
 - As a consequence of the statistically significant survival benefit, patients who were still receiving study treatment on the placebo arm were offered crossover to the pertuzumab arm.
 - The final exploratory analysis of OS is event driven and will take place when 385 events have been reached.
- No new safety signals compared to the primary analysis were reported with 1 more year of follow-up.
- These results indicate that combined HER2 blockade and chemotherapy with pertuzumab/trastuzumab/docetaxel can be considered a standard first-line therapy for patients with HER2-positive mBC.

Swain SM et al. *Proc SABCS 2012*;Abstract P5-18-26.

Research
To Practice®

Investigator Commentary: Confirmatory OS Analysis of CLEOPATRA – Pertuzumab with Trastuzumab and Docetaxel as First-Line Therapy for HER2-Positive mBC

Changes in the first-line setting for patients with mBC have been coming along rapidly. The primary analysis of CLEOPATRA reported a significant improvement in PFS, and, as expected, this current analysis reported an improvement in OS. The authors confirmed little increase in toxicity compared to dual taxane/trastuzumab therapy. This report solidifies that the appropriate strategy is to administer a taxane in combination with trastuzumab and pertuzumab in the first-line setting.

Recent data have shown similar efficacy with docetaxel/trastuzumab and vinorelbine/trastuzumab but lower toxicity with the vinorelbine versus the docetaxel combination. The results of CLEOPATRA have given us impetus for an ongoing trial called VELVET that I have the honor of leading. This Phase II trial is for patients who are eligible to receive first-line therapy for HER2-positive mBC. In the first cohort, which we have already enrolled, patients will receive pertuzumab and trastuzumab sequentially. The second cohort will receive pertuzumab and trastuzumab together. Vinorelbine will be given to both cohorts.

Interview with Edith A Perez, MD, January 17, 2013

Research
To Practice®